

Technical University of Denmark



Electron Microscopy for Morphological Analysis of Black Carbon Accumulated in Glaciers

Mateiu, Ramona Valentina; Nagatsuka, Naoko ; Goto-Azuma, Kumiko ; Wagner, Jakob Birkedal

Published in:

Book of Abstracts. DTU's Sustain Conference 2015

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Mateiu, R. V., Nagatsuka, N., Goto-Azuma, K., & Wagner, J. B. (2015). Electron Microscopy for Morphological Analysis of Black Carbon Accumulated in Glaciers. In Book of Abstracts. DTU's Sustain Conference 2015 [S-12] Lyngby: Technical University of Denmark (DTU).

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Electron Microscopy for Morphological Analysis of Black Carbon Accumulated in Glaciers

Ramona Valentina Mateiu^{1*}, Naoko Nagatsuka², Kumiko Goto-Azuma² and Jakob Birkedal Wagner¹

1: Technical University of Denmark, Copenhagen, Denmark; 2: National Institute of Polar Research, Tokyo, Japan

*Corresponding author email: ramona.mateiu@cen.dtu.dk

Since the mid-20th century the decrease in anthropogenic emission of Black Carbon in Europe and North America has been recorded as a decreased Black Carbon accumulation in Greenland. However, the present-day concentrations of Black Carbon in Greenland are higher compared to the Black Carbon concentrations from the pre-industrial time, and the Black Carbon concentration in the Arctic.

Accumulation of Black Carbon in the glaciers correlates with a decrease in the surface albedo and therefore Black Carbon is thought to be one of the causes that lead to the melting of the glaciers. The morphology of Black Carbon depends on its origin, the various processes during transportation and the incorporation process into snow.

In the present study we evaluate the use of electron microscopy for morphological analysis of Black Carbon. We use both scanning electron microscopy (SEM) and transmission electron microscopy (TEM) for the analysis of Black Carbon in snow collected from Greenland and Alaska. We show that there is a morphological preference for the Black Carbon depending on the snow origin: Black Carbon found in snow from Greenland has fewer monomers arranged in a lacy, chain like, shape, while Black Carbon found in snow from Alaska has a large number of monomers arranged in a compact, onion like, shape.

These preliminary results show that electron microscopy can be used for morphological analysis of Black Carbon accumulated in snow and glaciers, and that the Black Carbon particles present morphological differences depending on their origin.